arXiv:1107.3918v4 [hep-th] 18 Aug 2011

CQUEST 2011-0459 July 2011

$SU(N)$ quantum Racah coefficients & non-torus links

Zodinmawia[1] and P. Ramadevi[1,2]
[1] *Department of Physics,*
Indian Institute of Technology Bombay,
Mumbai 400 076, India
and
[2] *Center for Quantum Spacetime,*
Sogang University, Seoul, S.Korea

Abstract

It is well-known that the $SU(2)$ quantum Racah coefficients or the Wigner $6j$ symbols have a closed form expression which enables the evaluation of any knot or link polynomials in $SU(2)$ Chern-Simons field theory. Using isotopy equivalence of $SU(N)$ Chern-Simons functional integrals over three balls with one or more S^2 boundaries with punctures, we obtain identities to be satisfied by the $SU(N)$ quantum Racah coefficients. This enables evaluation of the coefficients for a class of $SU(N)$ representations. Using these coefficients, we can compute the polynomials for some non-torus knots and two-component links. These results are useful for verifying conjectures in topological string theory.

1 Introduction

Following the seminal work of Witten[1] on Chern-Simons theory as a theory of knots and links, generalised invariants[2, 3] for any knot or link can be directly obtained without going through the recursive procedure. For torus links, which can be wrapped on a two-torus T^2, using the torus link operators [4], explicit polynomial form of these invariants could be obtained. However for non-torus links, the generalised invariants [2] in $SU(N)$ Chern-Simons theory involves $SU(N)$ quantum Racah coefficients which are not known in closed form as known for $SU(2)$[5, 16]. This prevents in writing the polynomial form for the non-torus links.

For simple $SU(N)$ representations placed on knots, whose Young Tableau are $\square$, $\square\square$ and $\begin{smallmatrix}\square\\\square\end{smallmatrix}$, we had obtained some Racah coefficients from isotopy equivalence of knots and links which was useful to obtain polynomial for few non-torus knots[2, 3]. Going beyond these simple representations and finding the quantum Racah coefficients appeared to be a formidable task. In fact, determining these coefficients would help in verifying the topological string conjectures for a general non-torus knot or link observable proposed by Ooguri-Vafa[6, 7]. Using the few Racah coefficients data[2], Ooguri-Vafa conjecture for $4_1, 6_1$ non-torus knots as indicated in Fig. 1 were verified[8]. For verifying Ooguri-Vafa conjecture for the non-torus two-component links[7], we need to evaluate the non-torus link whose component knots carry different representations.

These non-torus links invariants will also be useful to generalise some of the results of recent papers[9, 10, 11, 12] where torus knots and links are studied. So, it is very important to determine the $SU(N)$ quantum Racah coefficients.

Using the correspondence between Chern-Simons functional integral and the correlator conformal blocks states in the corresponding Wess-Zumino conformal field theory[1], we can derive identities to be obeyed by the $SU(N)$ quantum Racah coefficients. Using the identities and the the properties of quantum dimensions for any N, we could determine the form of these coefficients for some class of $SU(N)$ representations. These coefficients are needed to obtain polynomial invariants of some non-torus knots and non-torus two-component links.

The plan of the paper is as follows: In section 2, we briefly review Chern-Simons functional integrals and properties of the Racah coefficients. In section 3, we systematically study the equivalence of states and obtain identities to be obeyed by the $SU(N)$ quantum Racah coefficients. In section 4, we tabulate the $SU(N)$ quantum Racah coefficients. In section 5, we use these coefficients to obtain the generalised polynomial invariant for a non-torus knot and a non-torus link and verify Ooguri-Vafa conjecture. Finally, we conclude with summary and discussions.

2 Chern-Simons Field Theory

Chern-Simons fields on S^3 with $U(1) \times SU(N)$ gauge group with levels k_1, k respectively is given by by the following action:

$$S = \frac{k_1}{4\pi} \int_{S^3} B \wedge dB + \frac{k}{4\pi} \int_{S^3} Tr \left(A \wedge dA + \frac{2}{3} A \wedge A \wedge A \right) , \qquad (2.1)$$

where B is the $U(1)$ gauge connection and A is the $SU(N)$ matrix valued gauge connection. The observables in this theory are Wilson loop operators:

$$W_{(R_1,n_1),(R_2,n_2),\ldots(R_r,n_r)}[\mathcal{L}] = \prod_{\beta=1}^{r} Tr_{R_\beta} U^A[\mathcal{K}_\beta] Tr_{n_\beta} U^B[\mathcal{K}_\beta] , \qquad (2.2)$$

where the holonomy of the gauge field A around a component knot $\mathcal{K}_\beta$, carrying a representation R_β, of a r-component link is denoted by $U^A[\mathcal{K}_\beta] = P[\exp \oint_{\mathcal{K}_\beta} A]$ and n_β is the $U(1)$ charge carried by the component knot $\mathcal{K}_\beta$. The expectation value of these Wilson loop operators are the link invariants:

$$\begin{aligned} V^{\{U(N)\}}_{(R_1,n_1),\ldots}[\mathcal{L}](q,\lambda) &= \langle W_{(R_1,n_1),\ldots}[\mathcal{L}] \rangle = \frac{\int [\mathcal{D}B][\mathcal{D}A] e^{iS} W_{(R_1,n_1),\ldots(R_r,n_r)}[\mathcal{L}]}{\int [\mathcal{D}B][\mathcal{D}A] e^{iS}} \\ &= V^{\{SU(N)\}}_{R_1,R_2,\ldots R_r}[\mathcal{L}] V^{\{U(1\}}_{n_1,n_2,\ldots,n_r}[\mathcal{L}] \ . \qquad (2.3) \end{aligned}$$



$(b_2^{(+)})^{(-1)}(b_1^{(-)})^{(-1)}(b_2^{(-)})^2$

4_1



$(b_2^{(+)})^{-1}(b_1^{(-)})^{-1}(b_3^{(-)})^{-1}(b_2^{(+)})^{-2}$

5_2



$(b_2^{(+)})^{-1}(b_1^{(-)})^{-3}(b_2^{(-)})^2$

6_1



$(b_2^{(-)})^{-1}(b_1^{(+)})^{-1}(b_3^{(+)})^{-1}(b_2^{(-)})^{-1}(b_3^{(-)})b_2^{(+)}$

6_2



$b_2^{(-)}b_1^{(+)}b_2^{(-)}(b_3^{(-)})^{-1}(b_2^{(+)})^{-2}$

6_3



$(b_2^{(+)})^{-1}(b_1^{(-)})^{-2}(b_3^{(-)})^{-2}(b_2^{(+)})^{-2}$

7_2



$b_2^{(-)}(b_1^{(+)})^2b_3^{(+)}(b_2^{(-)})^3$

7_3



$b_2^{(+)}b_1^{(-)}b_3^{(-)}b_2^{(+)}b_1^{(-)}b_3^{(-)}b_2^{(+)}$

7_4



$(b_2^{(-)})^{-1}(b_1^{(+)})^{-1}(b_3^{(+)})^{-1}(b_2^{(-)})^{-2}(b_3^{(+)})^{-1}(b_2^{(-)})^{-1}$

7_5



$(b_2^{(-)})^{-1}(b_1^{(+)})^{-1}(b_2^{(-)})^{-1}(b_3^{(-)})^2(b_2^{(-)})^{-2}$

7_6



$b_1^{(+)}b_1^{(-)}(b_2^{(-)})^{-1}(b_3^{(+)})^{-1}(b_2^{(-)})^{-1}b_1^{(-)}b_2^{(+)}$

7_7

Figure 1: Plat representation of non-torus knots upto 7-crossings



$(b_2^{(-)})^{-1} b_4^{(+)} (b_3^{(+)})^{-1} (b_2^{(-)})^{-1} b_4^{(+)}$ $(b_2^{(+)})^{-1} (b_1^{(-)})^{-1} (b_3^{(-)})^{-1} (b_2^{(+)})^{-3}$ $b_2^{(+)} b_1^{(-)} (b_2^{(-)})^{-2} b_3^{(-)} b_2^{(+)}$

5_1 6_2 6_3



$(b_2^{(-)})^{-1} (b_1^{(+)})^{-3} (b_2^{(-)})^{-1} b_3^{(-)} b_2^{(+)}$ $b_2^{(-)} b_1^{(+)} b_3^{(+)} b_2^{(-)} (b_3^{(-)})^{-1} (b_2^{(+)})^{-2}$ $(b_2^{(-)})^{-1} (b_1^{(+)})^{-1} (b_2^{(-)})^{-3} b_3^{(-)} b_2^{(+)}$

7_1 7_2 7_3

Figure 2: Plat representation of non-torus links upto 7-crossings

We make a specific choice of the U(1) charges and the coupling k_1 so that the above invariant is a polynomial in two variables $q = \exp\left(\frac{2\pi i}{k+N}\right)$, $\lambda = q^N$[13, 14]:

$$n_\beta = \frac{l_\beta}{\sqrt{N}} \ ; \ k_1 = k + N \ , \tag{2.4}$$

where l_β is the total number of boxes in the Young Tableau representation R_β. The $U(1)$ link invariant for the link with this substitution gives

$$V^{\{U(1)\}}_{\frac{l_1}{\sqrt{N}},\ldots,\frac{l_r}{\sqrt{N}}}[\mathcal{L}] = (-1)^{\sum_\beta l_\beta p_\beta} \exp\left(\frac{i\pi}{k+N}\sum_{\beta=1}^{r}\frac{l_\beta^2 p_\beta}{N}\right)\exp\left(\frac{i\pi}{k+N}\sum_{\alpha\neq\beta}\frac{l_\alpha l_\beta lk_{\alpha\beta}}{N}\right) \ , \tag{2.5}$$

where p_β is the framing number of the component knot $\mathcal{K}_\beta$ and $lk_{\alpha\beta}$ is the linking number between the component knots $\mathcal{K}_\alpha$ and $\mathcal{K}_\beta$. In order to directly evaluate $SU(N)$ link invariants, we need to use the following two ingredients:

1. The relation between $SU(N)$ Chern-Simons functional integral on the three-dimensional ball to the two-dimensional $SU(N)_k$ Wess-Zumino conformal field theory on the boundary of the three-ball[1].

2. Any knot or link can be drawn as a platting of braids[15].

In Fig. 1 and Fig. 2 we have drawn some non-torus knots and non-torus links as a plat representation of braids. We have labelled them in the Thistlewaithe notation and written their braid words. We have indicated the orientation and labelled the representation R_i on the component knots in the link. Note that $b_i^{(-)}$ ($\{b_i^{(-)}\}^{-1}$) in the braid word denotes right-handed crossing (left-handed crossing) between i-strand and $i+1$-th strand which are anti-parallelly oriented. Similarly $b_j^{(+)}$ ($\{b_j^{(+)}\}^{-1}$) denotes right-handed crossing (left-handed crossing) between j and $j+1$-th strand which are parallelly oriented. The plat representation of these non-torus knots and non-torus links except link 5_4 involves braids with four-strands. Hence we can view these knots and links except link 5_4 in S^3 as gluing of three-balls with 4-punctured boundary as shown in Fig. 3. There are two three-balls B_1 and B_3 with one S^2 boundary. A three-ball denoted as B_2 in Fig. 3 with two S^2 boundaries with a braid $\mathcal{B}$ which can represent any of the braid words corresponding to the non-torus knots and links in Figs. 1&2. The gluing of the three-balls are along the oppositely oriented S^2 boundaries. The functional integral over the ball B_3 is



Figure 3: Link in S^3 viewed as gluing of three-balls with four-punctured S^2 boundaries

given by a state $|\chi_0^{(2)}\rangle$ where the superscript denotes the label of the S^2 boundary. The representation R_i indicate that the lines are going into the S^2 boundary of the three-ball and the conjugate representation denotes the lines going out of the S^2 boundary. The state corresponding to a functional integral on a three-ball with an oppositely oriented boundary is written in a dual space along with conjugating all the representations as illustrated for the ball B_1. The expectation value of the Wilson-loop operator gives the link invariant for a nontorus link L

$$V_{R_1,R_2}^{SU(N)}[L] = \langle \Psi_0^{(1)}|\mathcal{B}\nu^{(1),(2)}|\chi_0^{(2)}\rangle \ . \tag{2.6}$$

These invariants multiplied with the $U(1)$ invariant (2.5) are polynomials in two variables $q = \exp(2\pi i/(k+N))$ and $\lambda = q^N$. In order to see the polynomial form, we write these states on a four-punctured boundary in a suitable basis of four-point conformal block of the $SU(N)_k$ Wess-Zumino conformal field theory. There are two different four-point conformal block bases as shown in Fig. 4 where $t \in (R_1 \otimes R_2) \cap (R_3 \times R_4)$ and $s \in (R_2 \otimes \bar{R}_3) \cap (\bar{R}_1 \times R_4)$. Using these bases, the states corresponding to three balls B_1, B_2 and B_3 in Fig. 3, can be expanded as



$\phi_t(R_1, R_2.\bar{R}_3, \bar{R}_4)$ $\hat{\phi}_s(R_1, R_2.\bar{R}_3, \bar{R}_4)$

Figure 4: four-point conformal block bases states

[2, 3]

$$\begin{aligned} \langle \Psi_0^{(1)}| &= \sqrt{dim_q R_1 dim_q R_2}\langle \phi_0(\bar{R}_1, R_1, R_2, \bar{R}_2)^{(1)}| && (2.7)\\ &= \sum_{s \in R_1 \otimes R_2} \epsilon_s^{(R_1,R_2)} \sqrt{dim_q s}\langle \hat{\phi}_s(\bar{R}_1, R_1, R_2, \bar{R}_2)^{(1)}| \\ \mathcal{B}\nu^{(1),(2)} &= \mathcal{B}|\phi_t(R_2, \bar{R}_2, \bar{R}_1, R_1)^{(1)}\rangle\langle \phi_t(\bar{R}_2, R_2, R_1, \bar{R}_1)^{(2)}| \\ &= \mathcal{B}|\hat{\phi}_s(R_2, \bar{R}_2, \bar{R}_1, R_1)^{(1)}\rangle\langle \hat{\phi}_s(\bar{R}_2, R_2, R_1, \bar{R}_1)^{(2)}| && (2.8)\\ |\chi_0^{(2)}\rangle &= \sqrt{dim_q R_1 dim_q R_2}|\phi_0(R_2, \bar{R}_2, \bar{R}_1, R_1)^{(2)}\rangle && (2.9)\\ &= \sum_{s \in R_1 \otimes R_2} \epsilon_s^{(R_1,R_2)} \sqrt{dim_q s}|\hat{\phi}_s(R_2, \bar{R}_2, \bar{R}_1, R_1)^{(2)}\rangle \ , && (2.10) \end{aligned}$$

where the subscript '0' in the basis state in eqns.(2.7, 2.9) denotes the singlet state. The states are written such that the invariant of a simple circle (also called unknot) carrying representation R is normalised as $dim_q R$, which is the quantum dimension of the representation R, defined as

$$dim_q R \ = \ \prod_{\alpha > 0} \frac{[\alpha.(\rho + \Lambda_R)]}{[\alpha.\rho]} \ , \tag{2.11}$$

where Λ_R denotes the highest weight of the representation R, α's are the positive roots and ρ is equal to the sum of the fundamental weights of the Lie group $SU(N)$. The square bracket refers to quantum number defined as

$$[n] = \frac{q^{\frac{n}{2}} - q^{-\frac{n}{2}}}{q^{\frac{1}{2}} - q^{-\frac{1}{2}}} \ . \tag{2.12}$$

The symbol $\epsilon_s^{(R_1,R_2)} = \pm 1$ which we will fix from equivalence of states in the next section. To operate the braid word $\mathcal{B}$ in eqn.(2.8), we need to find the eigenbasis of the braiding generators $b_i^{(\pm)}$'s.

The conformal block $|\phi_t(R_1, R_2, \bar{R}_3, \bar{R}_4)\rangle$ is suitable for the braiding operator $b_1^{(\pm)}$ and $b_3^{(\pm)}$. Similarly braiding in the middle two strands involving the operator $b_2^{(\pm)}$ requires the conformal block $|\hat{\phi}_s(R_1, R_2, \bar{R}_3, \bar{R}_4)\rangle$. That is,

$$\begin{aligned}
b_2^{(\pm)}|\hat{\phi}_s(R_1, R_2, \bar{R}_3, \bar{R}_4)\rangle &= \lambda_s^{(\pm)}(R_2, \bar{R}_3)|\hat{\phi}_s(R_1, \bar{R}_3, R_2, \bar{R}_4)\rangle \ , \\
b_1^{(\pm)}|\phi_t(R_1, R_2, \bar{R}_3, \bar{R}_4)\rangle &= \lambda_t^{(\pm)}(R_1, R_2)|\hat{\phi}_s(R_2, R_1, \bar{R}_3, \bar{R}_4)\rangle \ , \\
b_3^{(\pm)}|\hat{\phi}_s(R_1, R_2, \bar{R}_3, \bar{R}_4)\rangle &= \lambda_t^{(\pm)}(\bar{R}_3, \bar{R}_4)|\hat{\phi}_s(R_1, R_2, \bar{R}_4, \bar{R}_3)\rangle \ , \qquad (2.13)
\end{aligned}$$

where braiding eigenvalues $\lambda_t^{(\pm)}(R_1, R_2)$ in vertical framing are

$$\lambda_t^{(\pm)}(R_1, R_2) = \epsilon_{t;R_1,R_2}^{(\pm)} \left(q^{\frac{C_{R_1}+C_{R_2}-C_{R_t}}{2}} \right)^{\pm 1} \ . \qquad (2.14)$$

In this framing, framing number p_β for the component knot is equal to writhe w of that component knot which is the difference between total number of left-handed crossings and total number of right-handed crossing. For example, torus knots $4_1, 5_2$ in Fig. 1 have writhe w equal to 0 and 5 respectively. The symbol $\epsilon_{t;R_1,R_2}^{(\pm)}$ is a sign which can be fixed by studying equivalence of states or equivalence of links which we shall elaborate for a class of representations in the next section and C_R denotes the quadratic casimir for the representation R given by

$$C_R = \kappa_R - \frac{l^2}{2N} \ , \ \kappa_R = \frac{1}{2}[Nl + l + \sum_i (l_i^2 - 2il_i)] \ , \qquad (2.15)$$

where l_i is the number of boxes in the i-th row of the Young Tableau representation R and l is the total number of boxes. The two bases in Fig. 4 are related by a duality matrix a as follows:

$$|\phi_t(R_1, R_2, \bar{R}_3, \bar{R}_4)\rangle = a_{ts} \begin{bmatrix} R_1 & R_2 \\ \bar{R}_3 & \bar{R}_4 \end{bmatrix} |\hat{\phi}_s(R_1, R_2, \bar{R}_3, \bar{R}_4)\rangle \ . \qquad (2.16)$$

From the definition of t,s, we can see that the duality matrix a obeys the following properties:

$$a_{ts} \begin{bmatrix} R_1 & R_2 \\ \bar{R}_3 & \bar{R}_4 \end{bmatrix} = a_{st} \begin{bmatrix} \bar{R}_3 & R_2 \\ R_1 & \bar{R}_4 \end{bmatrix} = a_{\bar{s}\bar{t}} \begin{bmatrix} R_1 & \bar{R}_4 \\ \bar{R}_3 & R_2 \end{bmatrix} = a_{\bar{t}\bar{s}} \begin{bmatrix} \bar{R}_3 & \bar{R}_4 \\ R_1 & R_2 \end{bmatrix} = a_{ts} \begin{bmatrix} R_3 & R_4 \\ \bar{R}_1 & \bar{R}_2 \end{bmatrix} . \qquad (2.17)$$

If one of the representation is singlet (denoted by 0), we see that the matrix elements are

$$a_{ts}\begin{bmatrix} R_1 & 0 \\ \bar{R}_3 & \bar{R}_4 \end{bmatrix} = \delta_{tR_1}\delta_{s\bar{R}_3}\ ,\ a_{ts}\begin{bmatrix} 0 & R_2 \\ \bar{R}_3 & \bar{R}_4 \end{bmatrix} = \delta_{tR_2}\delta_{sR_4}\ ,$$

$$a_{ts}\begin{bmatrix} R_1 & R_2 \\ 0 & \bar{R}_4 \end{bmatrix} = \delta_{tR_4}\delta_{sR_2}\ ,\ a_{ts}\begin{bmatrix} R_1 & R_2 \\ \bar{R}_3 & 0 \end{bmatrix} = \delta_{tR_3}\delta_{s\bar{R}_1}\ . \qquad (2.18)$$

From the procedure presented in this section, we can write the $U(N)$ invariants of non-torus knots and links as a product of $U(1)$ invariant times $SU(N)$ invariant (2.6). For example, non-torus knot 5_2 with framing $p = 5$, the invariant will be

$$\begin{aligned} V_R^{\{U(N)\}}[5_2;p] &= (-1)^{5l} q^{\frac{5l^2}{2N}} \sum_{s,t,s'} \epsilon_s^{R,R}\ \sqrt{dim_q s}\ \epsilon_{s'}^{R,R}\ \sqrt{dim_q s'} (\lambda_s^{(+)}(R,R))^{-1} \\ & \qquad a_{ts}\begin{bmatrix} \bar{R} & R \\ R & \bar{R} \end{bmatrix} (\lambda_t^{(-)}(\bar{R},R))^{-2} a_{ts'}\begin{bmatrix} R & \bar{R} \\ \bar{R} & R \end{bmatrix} (\lambda_{s'}^{(+)}(R,R))^{-2}, \end{aligned} \qquad (2.19)$$

where l is the total number of boxes in the Young Tableau representing R and we have indicated the framing number of the knot. We could add additional framing p_1 by multiplying these invariants by a framing factor as follows:

$$\begin{aligned} V_R^{\{U(N)\}}[K;(p+p_1)] &= (-1)^{lp_1} q^{\frac{p_1 l^2}{2N}} (\lambda_0^{(-)}(R,\bar{R}))^{-p_1} V_R^{\{U(N)\}}[K;p] \\ &= (-1)^{lp_1} q^{p_1\kappa_R} V_R^{(U(N))}[K;p]\ . \end{aligned} \qquad (2.20)$$

So, to obtain 5_2 invariant with zero framing, we have to take $p_1 = -5$. Similarly the invariant for link 6_2 with linking number $lk = 3$ with framing p_1, p_2 on the component knots will be

$$\begin{aligned} V_{(R_1,R_2)}^{\{U(N)\}}[6_2;p_1,p_2] &= \prod_{i=1}^{2}\{(-1)^{l_{R_i}p_i} q^{p_i\kappa_{R_i}}\} q^{\frac{3 l_{R_1} l_{R_2}}{N}} \sum_{s,t,s'} \epsilon_s^{R_1,R_2}\ \sqrt{dim_q s} \qquad (2.21) \\ & \qquad \epsilon_{s'}^{\bar{R}_1,\bar{R}_2}\ \sqrt{dim_q s'} (\lambda_s^{(+)}(R_1,R_2))^{-3} a_{ts}\begin{bmatrix} R_2 & \bar{R}_1 \\ \bar{R}_2 & R_1 \end{bmatrix} \\ & \qquad (\lambda_t^{(-)}(\bar{R}_1,R_2))^{-2} a_{ts'}\begin{bmatrix} \bar{R}_1 & R_2 \\ R_1 & \bar{R}_2 \end{bmatrix} (\lambda_{s'}^{(+)}(\bar{R}_1,\bar{R}_2))^{-1}\ , \end{aligned}$$

where l_{R_i} is the total number of boxes in the Young diagram representation R_i placed on the component knots in link 6_2. However to see the polynomial form of these link invariants, we have to determine the coefficients of the duality matrices. Unlike the $SU(2)$ duality matrices[5, 16], we do not have a closed form expression for SU(N) duality matrices.

In the following section, we will use equivalence of states to obtain the sign (2.14, 2.7) and also derive identities satisfied by the coefficients of the duality matrix. This enables the evaluation of duality matrix elements for some class of representations.

Once we have these coefficients, we could evaluate the framed link invariants and obtain the reformulated invariants[17]

$$\begin{aligned} f_{R_1,R_2,\ldots R_r}(q,\lambda) &= \sum_{d,m=1}^{\infty}(-1)^{m-1}\frac{\mu(d)}{dm}\sum_{\{\vec{k}^{(\alpha j)},R_{\alpha j}\}}\times \\ & \prod_{\alpha=1}^{r}\chi_{R_\alpha}\left(C\left((\sum_{j=1}^{m}\vec{k}^{(\alpha j)})_d\right)\right)\prod_{j=1}^{m}\frac{|C(\vec{k}^{(\alpha j)})|}{\ell_{\alpha j}!}\times \\ & \chi_{R_{\alpha j}}(C(\vec{k}^{(\alpha j)}))\ V^{(U(N))}_{R_{1j},R_{2j},\ldots R_{rj}}[L;\{p_\alpha\}](q^d,\lambda^d)\ , \quad (2.22) \end{aligned}$$

where $\mu(d)$ is the Moebius function defined as follows: if d has a prime decomposition $(\{p_i\})$, $d=\prod_{i=1}^{a}p_i^{m_i}$, then $\mu(d)=0$ if any of the m_i is greater than one. If all $m_i=1$, then $\mu(d)=(-1)^a$. The second sum in the above equation runs over all vectors $\vec{k}^{(\alpha j)}$, with $\alpha=1,\ldots r$ and $j=1,\ldots m$, such that $\sum_{\alpha=1}^{r}|\vec{k}^{(\alpha j)}|>0$ for any j and over representations $R_{\alpha j}$. Further $\vec{k}_d$ is defined as follows: $(\vec{k}_d)_{di}=k_i$ and has zero entries for the other components. Therefore, if $\vec{k}=(k_1,k_2,\ldots)$, then

$$\vec{k}_d=(0,\ldots,0,k_1,0,\ldots,0,k_2,0,\ldots), \quad (2.23)$$

where k_1 is in the d-the entry, k_2 in the $2d$-th entry, and so on. Here $C(\vec{k})$ denotes the conjugacy class determined by the sequence $(k_1,k_2,\cdots)$ (i.e there are k_1 1-cycles, k_2 2-cycles etc) in the permutation group S_ℓ ($\ell=\sum_j jk_j$). For a Young Tabeleau representation R with ℓ number of boxes, $\chi_R(C(\vec{k}))$ gives the character of the conjugacy class $C(\vec{k})$ in the representation R. The reformulated invariants for r-component links are expected to obey Ooguri-Vafa conjecture[7]:

$$f_{(R_1,R_2,\ldots R_r)}(q,\lambda)=(q^{1/2}-q^{-1/2})^{r-2}\sum_{Q,s}N_{(R_1,\ldots R_r),Q,s}q^s\lambda^Q\ , \quad (2.24)$$

where $N_{(R_1,\ldots R_r),Q,s}$ are integers. After determining the identities and some of the matrix elements of the duality matrices in the following two sections, we will obtain the polynomial invariant for 5_2 knot and 6_2 link and verify eqn.(2.24) in section 5.

3 Duality Matrix identities

We had elaborated in the previous section on writing states (2.7,2.8,2.9) corresponding to Chern-Simons functional integral on three balls. We can determine the following coefficients of the duality matrix by comparing eqn.(2.16) and eqns.(2.7,2.9):

$$a_{0s}\begin{bmatrix}\bar{R}_1 & R_1\\ R_2 & \bar{R}_2\end{bmatrix} = \epsilon_s^{(R_1,R_2)}\frac{\sqrt{dim_q s}}{\sqrt{dim_q R_1 dim_q R_2}}~. \tag{3.1}$$

This relation along with the property (2.18) suggests that

$$a_{ts}\begin{bmatrix}R_1 & R_2\\ \bar{R}_3 & \bar{R}_4\end{bmatrix} = \epsilon_{R_1}\epsilon_{R_2}\epsilon_{\bar{R}_3}\epsilon_{\bar{R}_4}\sqrt{dim_q s}\sqrt{dim_q t}\begin{Bmatrix}R_1 & R_2 & t\\ \bar{R}_3 & \bar{R}_4 & s\end{Bmatrix}~, \tag{3.2}$$

where $\epsilon_{R_i} = \pm 1 = \epsilon_{\bar{R}_i}$ and $\epsilon_0 = 1$. The term in parenthesis is similar to the $SU(2)$ quantum Wigner $6j$ symbol but requires appropriate conjugation of representations under interchange of columns in the following way:

$$\begin{Bmatrix}R_1 & R_2 & t\\ \bar{R}_3 & \bar{R}_4 & s\end{Bmatrix} = \begin{Bmatrix}t & \bar{R}_2 & R_1\\ s & \bar{R}_4 & \bar{R}_3\end{Bmatrix} = \begin{Bmatrix}\bar{R}_1 & t & R_2\\ \bar{R}_3 & \bar{s} & R_4\end{Bmatrix}~. \tag{3.3}$$

Using (2.18) and the relation to quantum Wigner symbol with the above properties, the $SU(N)$ duality matrix can be called as SU(N) quantum Racah coefficients and hence we propose that the coefficients obey the following property:

$$a_{ts}\begin{bmatrix}R_1 & R_2\\ \bar{R}_3 & \bar{R}_4\end{bmatrix} = \frac{\sqrt{dim_q t dim_q s}}{\sqrt{dim_q R_1 dim_q R_3}}\epsilon_{R_1}\epsilon_{\bar{R}_3}(\epsilon_s\epsilon_t)^{-1}a_{R_1\bar{R}_3}\begin{bmatrix}t & \bar{R}_2\\ s & \bar{R}_4\end{bmatrix}~. \tag{3.4}$$

Using this property, we can relate the sign in eqn.(3.1) as

$$\epsilon_s^{(R_1,R_2)} = \epsilon_{R_1}\epsilon_{R_2}(\epsilon_s)^{-1}~. \tag{3.5}$$

Starting from the state $\nu^{(1),(2)}$ in eqn.(2.8) and the duality relation (2.16), we observe that the Racah coefficients must obey the following identities:

$$\sum_s a_{ts} \begin{bmatrix} R_1 & \bar{R}_2 \\ \bar{R}_3 & \bar{R}_4 \end{bmatrix} a_{t's} \begin{bmatrix} R_1 & \bar{R}_2 \\ \bar{R}_3 & \bar{R}_4 \end{bmatrix} = \delta_{tt'} \ , \tag{3.6}$$

$$\sum_t a_{ts} \begin{bmatrix} R_1 & \bar{R}_2 \\ \bar{R}_3 & \bar{R}_4 \end{bmatrix} a_{ts'} \begin{bmatrix} R_1 & \bar{R}_2 \\ \bar{R}_3 & \bar{R}_4 \end{bmatrix} = \delta_{ss'} \ . \tag{3.7}$$

3.1 Fixing signs of the braiding eigenvalues



Figure 5: Unknot drawn in two equivalent ways

From Fig. 5, we can write the invariant for the unknot in two equivalent ways giving the following constraint equation:

$$\sum_s dim_q s \lambda_s^{(+)}(R,R) = \lambda_0^{(-)}(R,\bar{R}) dim_q R \ . \tag{3.8}$$

Taking $\epsilon^{(-)}_{0;R,\bar{R}} = 1$, we can determine the signs $\epsilon^{(+)}_{s;R,R}$ which satisfies the above equation. We can write a general form for the sign for a class of representations $R_n = \overset{\longleftarrow n \longrightarrow}{\square\square\square\square\square\square}$, the irreducible representations $\rho_\ell \in R_n \otimes R_n$ in $SU(N)_k$ Wess-Zumino conformal field theory in the large k limit will be



.

The sign $\epsilon^{(+)}_{\rho_\ell;R_n,R_n} = (-1)^{(n-\ell)}$. Similarly, for antiparallely oriented strands, the irreducible representations in $\tilde{\rho}_\ell \in R_n \otimes \bar{R}_n$ are



Here boxes with dot represents a column of length $N-1$. We take the sign $\epsilon^{(-)}_{\tilde{\rho}_\ell;R_n,\bar{R}_n} = (-1)^\ell$ which is $+1$ for the singlet $\ell = 0$. We can generalise these results for tensor product of two different symmetric representations $\rho_\ell \in R_n \otimes R_m$ and $\tilde{\rho}_\ell \in R_n \otimes \bar{R}_m$ as

$$\epsilon^{(+)}_{\rho_\ell}(R_n, R_m) = (-1)^{\frac{n+m}{2}-\ell} \ ; \ \epsilon^{(-)}_{\tilde{\rho}_\ell}(R_n, \bar{R}_m) = (-1)^{\frac{n-m}{2}-\ell} \ , \tag{3.9}$$

where we assume $n \geq m$ and $\ell = (n-m)/2, (n-m)/2+1, \ldots (n+m)/2$. Similarly for antisymmetric representations $\hat{R}_n$ placed on antiparallely oriented strands, the irreducible representations $\tilde{\hat{\rho}}_\ell \in \hat{R}_n \otimes \bar{\hat{R}}_n$ are



.

For this case, the sign $\epsilon^{(-)}_{\tilde{\hat{\rho}}_\ell;\hat{R}_n,\bar{\hat{R}}_n} = (-1)^\ell$. If we replace $N-n = n$ in the above tensor product, we get irreducible representations $\hat{\rho}_\ell \in \hat{R}_n \otimes \hat{R}_n$ for parallelly oriented strands carrying antisymmetric representation whose sign will be $\epsilon^{(+)}_{\hat{\rho}_\ell;\hat{R}_n,\hat{R}_n} = (-1)^{2n-\ell}$. The signs for antisymmetric representations can be similarly generalised for $\hat{\rho}_\ell \in \hat{R}_n \otimes \hat{R}_m$ and $\tilde{\hat{\rho}}_\ell \in \hat{R}_n \otimes \bar{\hat{R}}_m$ as

$$\epsilon^{(+)}_{\hat{\rho}_\ell;\hat{R}_n,\hat{R}_m} = (-1)^{n+m-\ell} \ ; \ \epsilon^{(-)}_{\tilde{\hat{\rho}}_\ell;\hat{R}_n,\bar{\hat{R}}_m} = (-1)^{n-m-\ell} \ , \tag{3.10}$$

where $n \geq m \& n+m \leq N$ and $\ell = 0, 1, \ldots m$ for parallel strands. Similarly for antiparallel strands with $N-m \geq n$, $\ell = n-m, n-m+1, \ldots n$.

It is possible to fix the signs for the mixed representation but cannot be written in the most general form as done for the symmetric and the antisymmetric representations. Some of the mixed representation signs are given in the earlier papers[8, 18]. For simplicity, we will confine to the symmetric or the antisymmetric representations placed on the component knots with the defined signs which will be useful for writing the Racah coefficients. In the following subsection, we will study equivalence of states which is needed to obtain the Racah coefficients.

3.2 More identities of Racah coefficients from equivalence of states

We can view the two three balls in Fig.6 as two equivalent states: The three-ball


oppositely oriented S^2 boundary
1
R_2
$\bar{R}_1$
$\equiv$
1
2
2
$\mathcal{B} = (b_2^{(+)})^{-1} b_3^{(-)}$
R_2
$\bar{R}_1$
R_1
$|\hat{\Psi}_0^{(1)}\rangle$
$\mathcal{B}\nu^{(1),(2)}$
$|\chi_0^{(2)}\rangle$

Figure 6: Two equivalent three-balls

corresponding to the state $|\hat{\Psi_0}^{(1)}\rangle$ can be glued with a similar three-ball with oppositely oriented S^2 boundary to give two unknots. So, this state can be represented as

$$|\hat{\Psi_0}^{(1)}\rangle = \epsilon(R_1, R_2)\sqrt{dim_q R_1 dim_q R_2}|\hat{\phi}_0(R_2\bar{R}_1 R_1 \bar{R}_2)^{(1)}\rangle \tag{3.11}$$

$$\begin{aligned} &= (b_2^{(+)})^{-1} b_3^{(-)} \nu^{(1),(2)} |\chi_0^{(2)}\rangle \\ &= \sum_s a_{0s} \begin{bmatrix} R_2 & \bar{R}_2 \\ \bar{R}_1 & R_1 \end{bmatrix} (\lambda_s^{(+)}(R_1, R_2))^{-1} a_{ts} \begin{bmatrix} R_2 & \bar{R}_1 \\ \bar{R}_2 & R_1 \end{bmatrix} \\ &\quad \times \lambda_t^{(-)}(R_2, \bar{R}_1)|\hat{\phi}_t(R_2\bar{R}_1 R_1\bar{R}_2)^{(1)}\rangle \ . \end{aligned} \tag{3.12}$$

From eqns.(3.11,3.12) and similar relations for braid word $\mathcal{B} = b_2^{(+)}(b_3^{(-)})^{-1}$, we can deduce the following identity

$$\sum_s a_{0s}\begin{bmatrix} R_2 & \bar{R}_2 \\ \bar{R}_1 & R_1 \end{bmatrix} (\lambda_s^{(+)}(R_1,R_2))^{\mp 1} a_{ts}\begin{bmatrix} R_2 & \bar{R}_1 \\ \bar{R}_2 & R_1 \end{bmatrix} = \epsilon(R_1,R_2)(\lambda_t^{(-)}(R_2,\bar{R}_1))^{\mp 1} a_{t0}\begin{bmatrix} R_2 & \bar{R}_1 \\ R_1 & \bar{R}_2 \end{bmatrix}. \qquad (3.13)$$

Using the data from $SU(2)$ [16], we can fix the signs $\epsilon(R_1,R_2)$ and the signs in the duality matrix for the class of symmetric or antisymmetric representations. Suppose we take symmetric representations for R_1, R_2 then the sign $\epsilon(R_n,R_m) = (-1)^{min(n,m)}$ and the signs in the duality matrix is $\epsilon_{R_n} = (-1)^{n/2}$. Similarly, for antisymmetric representations for R_1, R_2, $\epsilon(\hat{R}_n,\hat{R}_m) = (-1)^{min(2n,2m)}$ and $\epsilon_{\hat{R}_n} = (-1)^n$. With this sign convention, the above identity enables fixing some of the coefficients of the duality matrix which we tabulate in the next section.

The well-known braiding identity relates the two three-balls with two S^2 boundaries as pictorially shown in Fig. 7. Operating these braiding operators on the two S^2 boundary states, we can obtain the following identity for the duality matrix:



Figure 7: Braiding Identity

$$\sum_{s,t} \lambda_s^{(+)}(R_1,R_2) a_{st}\begin{bmatrix} R_2 & R_1 \\ \bar{R}_3 & \bar{R}_4 \end{bmatrix} \{\lambda_t^{(-)}(R_1,\bar{R}_3)\}^{-1} a_{\tilde{s}t}\begin{bmatrix} R_2 & \bar{R}_3 \\ R_1 & \bar{R}_4 \end{bmatrix} \{\lambda_{\tilde{s}}^{(-)}(R_2,\bar{R}_3)\}^{-1}$$

$$= \sum_{s,t,s',t'} a_{st}\begin{bmatrix} R_1 & R_2 \\ \bar{R}_3 & \bar{R}_4 \end{bmatrix} \{\lambda_t^{(-)}(R_2,\bar{R}_3)\}^{-1} a_{s't}\begin{bmatrix} R_1 & \bar{R}_3 \\ R_2 & \bar{R}_4 \end{bmatrix} \{\lambda_{s'}^{(-)}(R_1,\bar{R}_3)\}^{-1}$$

$$\times\, a_{s't'}\begin{bmatrix} \bar{R}_3 & R_1 \\ R_2 & \bar{R}_4 \end{bmatrix} \lambda_{t'}^{(+)}(R_1,R_2) a_{\tilde{s}t'}\begin{bmatrix} \bar{R}_3 & R_2 \\ R_1 & \bar{R}_4 \end{bmatrix}. \qquad (3.14)$$

3.2.1 Six Punctured S^2 boundaries

We have obtained these identities by studying equivalence of three-balls with four-punctured S^2 boundaries. We shall now look at a generalisation of Fig .6 for three-balls with six-punctured S^2 boundaries as depicted in Fig. 8 where the braiding operator $\mathcal{B} = g_6 = (\{b_4^{(+)}\}^{-1} b_3^{(-)})(\{b_5^{(+)}\}^{-1} b_4^{(-)} \{b_3^{(+)}\}^{-1} b_2^{(-)})$.

The Chern-Simons functional integral on these three-balls corresponds to a state in a space of six-point correlator conformal blocks in $SU(N)_k$ Wess-Zumino conformal field theory. They can be expanded in a convenient six point conformal block bases. Two such basis states are drawn in Fig .9. In terms of these six-point



Figure 8: Six-punctured boundary


R_1
R_2
R_3
R_4
R_5
R_6
s_1
s_2
s_3
$\phi_{s_1,s_2,s_3}(R_1, R_2, R_3, R_4, R_5, R_6)$
R_3
R_4
R_2
t_1
R_5
t_2
R_1
R_6
t_3
$\hat{\phi}_{t_1,t_2,t_3}(R_1, R_2, R_3, R_4, R_5, R_6)$

Figure 9: Six point conformal block bases

conformal block bases, we can relate the state $|\Psi\rangle$ and the state $|\xi\rangle$ as follows:

$$\begin{aligned}|\Psi\rangle &= \epsilon(R_1,R_2,R_3)\prod_{i=1}^{3}\sqrt{dim_q R_i}|\hat{\phi}_{0,0,0}(R_1,R_2,R_3,\bar{R}_3,\bar{R}_2,\bar{R}_1)\rangle = \mathcal{B}\nu^{(1),(2)}|\xi\rangle \\ &= \mathcal{B}\prod_{i=1}^{3}\sqrt{dim_q R_i}|\phi_{0,0,0}(R_1,\bar{R}_1,R_2,\bar{R}_2,R_3,\bar{R}_3)\rangle\ , \end{aligned} \tag{3.15}$$

where $\epsilon(R_1,R_2,R_3)=\pm 1=\epsilon(R_1,R_2)\epsilon(R_1,R_3)\epsilon(R_2,R_3)$ for symmetric and anti-symmetric representations. Applying the braiding operator $\mathcal{B}=g_6$ on the six-point conformal block in the above equation, we obtain the following relation:

$$\begin{aligned}\epsilon(R_1,R_2,R_3)|\hat{\phi}_{0,0,0}(R_1,R_2,R_3,\bar{R}_3,\bar{R}_2,\bar{R}_1)\rangle &= \sum \lambda^{(-)}_{q_1}(\bar{R}_1,R_2)\{\lambda^{(+)}_{s_1}(\bar{R}_1,\bar{R}_2)\}^{-1} \\ \times\lambda^{(-)}_{\mu_1}(\bar{R}_1,R_3)\{\lambda^{(+)}_{\nu_1}(\bar{R}_1,\bar{R}_3)\}^{-1}\lambda^{(-)}_{p_2}(\bar{R}_2,R_3)\{\lambda^{(+)}_{\mu_2}(\bar{R}_2,\bar{R}_3)\}^{-1} a_{t_1 0}&\begin{bmatrix}0 & R_2\\ \bar{R}_2 & 0\end{bmatrix} \\ \times a_{0q_1}\begin{bmatrix}R_1 & \bar{R}_1\\ R_2 & \bar{t}_1\end{bmatrix} a_{t_1 s_1}\begin{bmatrix}p_1 & \bar{R}_1\\ \bar{R}_2 & 0\end{bmatrix} a_{p_1 q_1}\begin{bmatrix}R_1 & R_2\\ \bar{R}_1 & \bar{t}_1\end{bmatrix} a_{t_2 s_1}&\begin{bmatrix}p_1 & \bar{R}_2\\ \bar{R}_1 & 0\end{bmatrix} a_{0\mu_1}\begin{bmatrix}t_2 & \bar{R}_1\\ R_3 & \bar{R}_3\end{bmatrix} \\ \times a_{t_2 p_2}\begin{bmatrix}p_1 & \bar{R}_2\\ R_3 & \bar{\nu}_1\end{bmatrix} a_{\nu_1\mu_1}\begin{bmatrix}t_2 & R_3\\ \bar{R}_1 & \bar{R}_3\end{bmatrix} a_{t_3 p_2}\begin{bmatrix}p_1 & R_3\\ \bar{R}_2 & \bar{\nu}_1\end{bmatrix} a_{\nu_1\mu_2}&\begin{bmatrix}t_3 & \bar{R}_2\\ \bar{R}_3 & \bar{R}_1\end{bmatrix} a_{\nu_2\mu_2}\begin{bmatrix}t_3 & \bar{R}_3\\ \bar{R}_2 & \bar{R}_1\end{bmatrix} \\ \times a_{t_3 p_3}\begin{bmatrix}p_1 & R_3\\ \bar{R}_3 & \bar{\nu}_2\end{bmatrix}|\phi_{p_1,p_3,\bar{\nu}_2}(R_1,R_2,R_3,\bar{R}_3,\bar{R}_2,\bar{R}_1)\rangle\ . \end{aligned} \tag{3.16}$$

We can simplify the RHS of the above expression using the property (2.18). Further, the summation over index q_1,μ_1 can be done using the identity (3.13). The simplified equation suggests another identity for the Racah coefficients:

$$\begin{aligned}&\sum_s a_{ts}\begin{bmatrix}R_1 & R_2\\ \bar{R}_3 & \bar{R}_4\end{bmatrix}\epsilon_s q^{\frac{\pm C_s}{2}} a_{ls}\begin{bmatrix}R_1 & \bar{R}_3\\ R_2 & \bar{R}_4\end{bmatrix} \\ &\qquad = (\epsilon_t\ \epsilon_l)^{-1} q^{\frac{\mp(C_t+C_l)}{2}}\prod_{i=1}^{4}(\epsilon_{R_i} q^{\frac{\pm(C_{R_i})}{2}}) a_{tl}\begin{bmatrix}R_2 & R_1\\ \bar{R}_3 & \bar{R}_4\end{bmatrix}\ . \end{aligned} \tag{3.17}$$

This is a generalisation of identity (3.13). Using this identity, we can do the summation over index p_2 and μ_2 and further simplify the RHS of the expression (3.16). The close similarity of these $SU(N)$ coefficients to the $SU(2)$ Racah coefficient identities suggests that all the identities of $SU(2)$ quantum coefficients must be generalisable to the $SU(N)$ coefficients and hence we postulate the following identity

$$\sum_{l_1} a_{r_2 l_1}\begin{bmatrix} r_1 & R_3 \\ R_4 & R_5 \end{bmatrix} a_{r_1 l_2}\begin{bmatrix} R_1 & R_2 \\ l_1 & R_5 \end{bmatrix} a_{\bar{l}_1 l_3}\begin{bmatrix} \bar{l}_2 & R_2 \\ R_3 & R_4 \end{bmatrix} = a_{r_2 l_2}\begin{bmatrix} R_1 & l_3 \\ R_4 & R_5 \end{bmatrix} a_{r_1 l_3}\begin{bmatrix} R_1 & R_2 \\ R_3 & \bar{r}_2 \end{bmatrix} , \qquad (3.18)$$

where we have appropriately chosen conjugate representations which are consistent with the definition of the Racah matrix. Again, using the above identity with $l_3 = 0$, we can do the the summation over ν_1 index in the simplified RHS of eqn.(3.16). Finally, we see the eqn.(3.16) reducing to

$$|\hat{\phi}_{0,0,0}(R_1, R_2, R_3, \bar{R}_3, \bar{R}_2, \bar{R}_1)\rangle = a_{p_1 0}\begin{bmatrix} R_1 & R_2 \\ \bar{R}_2 & \bar{R}_1 \end{bmatrix} |\phi_{p_1,0,p_1}(R_1, R_2, R_3, \bar{R}_3, \bar{R}_2, \bar{R}_1)\rangle \ . \qquad (3.19)$$

Using the properties of the duality matrix, RHS can be seen to be LHS. This elaborate exercise on the equivalence of two states corresponding to six-punctured boundary confirms that the correctness of the identity (3.18). Armed with these identities, we try to determine the Racah coefficients for some representations which we present in the next section.

4 $SU(N)$ quantum Racah coefficients

We shall use the properties and identities derived in the previous sections to obtain the duality matrix coefficients which will be useful for computing the non-torus knot and non-torus links.

For knots, all the strands carry same representation. So for obtaining non-torus knot invariants, we have to evaluate two types of Racah matrices-namely.,

$$a_{\hat{l}\hat{s}}\begin{bmatrix} R & \bar{R} \\ R & \bar{R} \end{bmatrix} \ , a_{l\hat{s}}\begin{bmatrix} R & R \\ \bar{R} & \bar{R} \end{bmatrix} \ , \qquad (4.1)$$

where first type can be shown to be a symmetric matrix from the properties of the Racah matrix. We could evaluate the symmetric first type Racah coefficients for

$R = \square, \square\square, \begin{smallmatrix}\square\\\square\end{smallmatrix}$ using eqns.(3.6,3.7) [2, 3]. However, for the second type Racah matrix we could only evaluate the coefficients for the fundamental representation($R = \square$).

Similarly for the two-component links, we can place two representations R_1, R_2 on the component knots. In this case, we can have three types of Racah matrices as follows:

$$a_{\hat{l}\hat{t}} \begin{bmatrix} R_1 & \bar{R}_1 \\ R_2 & \bar{R}_2 \end{bmatrix} \, , \; a_{l\hat{s}} \begin{bmatrix} R_1 & R_2 \\ \bar{R}_2 & \bar{R}_1 \end{bmatrix} \, , \; a_{l\hat{t}} \begin{bmatrix} R_1 & R_2 \\ \bar{R}_1 & \bar{R}_2 \end{bmatrix} \; . \tag{4.2}$$

Now, we will present the Racah coefficients for some representations which will be useful to compute the non-torus knot and link polynomials in $U(N)$ Chern-Simons theory.

4.1 Coefficients when R_1 is fundamental

1. For the simplest fundamental representation $R = \square$, the two types of Racah coefficient matrices are[2, 3]:

$$a_{ts} \left[\begin{array}{cc} R = \square & \bar{R} = \boxdot \\ R = \square & \bar{R} = \boxdot \end{array} \right] = \tfrac{1}{dim_q R} \left(\begin{array}{c|cc} & s = 0 & \boxdot\square \\ \hline t = 0 & -1 & \sqrt{[N-1][N+1]} \\ \boxdot\square & \sqrt{[N-1][N+1]} & 1 \end{array} \right) ,$$

$$a_{ts} \left[\begin{array}{cc} R = \square & R = \square \\ \bar{R} = \boxdot & \bar{R} = \boxdot \end{array} \right] = \tfrac{1}{dim_q R} \left(\begin{array}{c|cc} & s = 0 & \boxdot\square \\ \hline t = \begin{smallmatrix}\square\\\square\end{smallmatrix} & -\sqrt{\frac{[N][N-1]}{[2]}} & \sqrt{\frac{[N][N+1]}{[2]}} \\ \square\square & \sqrt{\frac{[N][N+1]}{[2]}} & \sqrt{\frac{[N][N-1]}{[2]}} \end{array} \right) ,$$

   where $dim_q(R = \square) = [N]$.

2. Next, we look at Racah coefficient matrices where $R_1 = \square \neq R_2$. This will be useful for the computation of two-component links.

$$a_{ts}\begin{bmatrix} R_1=\ydiagram{1} & \bar{R}_1=\begin{ytableau}\cdot\end{ytableau} \\ R_2=\ydiagram{2} & \bar{R}_2=\begin{ytableau}\cdot & \cdot\end{ytableau}\end{bmatrix}=\frac{1}{K}\left(\begin{array}{c|cc} & s=\begin{ytableau} \, & \, \\ \cdot \end{ytableau} & \begin{ytableau}\cdot & \, & \,\end{ytableau} \\ \hline t=0 & -\sqrt{[N]} & \sqrt{\frac{[N-1][N][N+2]}{[2]}} \\ \begin{ytableau}\cdot & \,\end{ytableau} & \sqrt{\frac{[N-1][N][N+2]}{[2]}} & \sqrt{[N]}\end{array}\right),$$

where $K=\sqrt{dim_q R_1 dim_q R_2}$. Similarly, the second and third type Racah matrix coefficients for $R_1=\ydiagram{1}, R_2=\ydiagram{2}$ are

$$a_{ts}\begin{bmatrix} R_1=\ydiagram{1} & R_2=\ydiagram{2} \\ \bar{R}_1=\begin{ytableau}\cdot\end{ytableau} & \bar{R}_2=\begin{ytableau}\cdot & \cdot\end{ytableau}\end{bmatrix}=\left(\begin{array}{c|cc} & s=\begin{ytableau} \, & \, \\ \cdot \end{ytableau} & \begin{ytableau}\cdot & \, & \,\end{ytableau} \\ \hline t=\ydiagram{2,1} & -\sqrt{\frac{[N-1]}{[3][N+1]}} & \sqrt{\frac{[2][N+2]}{[3][N+1]}} \\ \ydiagram{3} & \sqrt{\frac{[2][N+2]}{[3][N+1]}} & \sqrt{\frac{[N-1]}{[3][N+1]}}\end{array}\right),$$

$$a_{ts}\begin{bmatrix} R_1=\ydiagram{1} & R_2=\ydiagram{2} \\ \bar{R}_2=\begin{ytableau}\cdot & \cdot\end{ytableau} & \bar{R}_1=\begin{ytableau}\cdot\end{ytableau}\end{bmatrix}=\frac{1}{K}\left(\begin{array}{c|cc} & s=0 & \begin{ytableau}\cdot & \,\end{ytableau} \\ \hline t=\ydiagram{2,1} & -\sqrt{\frac{[N-1][N][N+1]}{[3]}} & \sqrt{\frac{[N][N+1][N+2]}{[2][3]}} \\ \ydiagram{3} & \sqrt{\frac{[N][N+1][N+2]}{[2][3]}} & \sqrt{\frac{[N-1][N][N+1]}{[3]}}\end{array}\right).$$

3. Interestingly, we could find the coefficients for $R_1=\ydiagram{1}, R_2=\overset{\xleftrightarrow{n}}{\ydiagram{5}}$ using the identities. The corresponding conjugate representations are $\bar{R}_1=\begin{ytableau}\cdot\end{ytableau}$, $\bar{R}_2=\overset{\xleftrightarrow{n}}{\begin{ytableau}\cdot & \cdot & \cdot & \cdot & \cdot\end{ytableau}}$ The three types of Racah coefficients for this class of representations are

$$a_{ts}\begin{bmatrix} R_1 & \bar{R}_1 \\ R_2 & \bar{R}_2\end{bmatrix}=\frac{1}{K}\left(\begin{array}{c|cc} \text{t} & s=\overset{\xleftrightarrow{n}}{\begin{ytableau} \, & \, & \, & \, & \, \\ \cdot \end{ytableau}} & \overset{\xleftrightarrow{n+1}}{\begin{ytableau}\cdot & \, & \, & \, & \, & \,\end{ytableau}} \\ \hline 0 & -\sqrt{\frac{[N][N+1]...[N+n-2]}{[n-1]!}} & \sqrt{\frac{[N-1][N]...[N+n]}{[N+n-1][n]!}} \\ \begin{ytableau}\cdot & \,\end{ytableau} & \sqrt{\frac{[N-1][N]...[N+n]}{[N+n-1][n]!}} & \sqrt{\frac{[N][N+1]...[N+n-2]}{[n-1]!}}\end{array}\right),$$

$$a_{ts}\begin{bmatrix} R_1 & R_2 \\ \bar{R}_1 & \bar{R}_2 \end{bmatrix} = \left(\begin{array}{c|cc} & s = \overset{\longleftarrow n \longrightarrow}{\ydiagram{5,1}} & \overset{\longleftarrow n+1 \longrightarrow}{\ydiagram{6}} \\ \hline t = \overset{\longleftarrow n \longrightarrow}{\ytableaushort{{}{}{}{}{},{\cdot}}} & -\sqrt{\frac{[N-1]}{[n+1][N+n-1]}} & \sqrt{\frac{[n][N+n]}{[n+1][N+n-1]}} \\ \overset{\longleftarrow n+1 \longrightarrow}{\ytableaushort{{\cdot}{}{}{}{}{}}} & \sqrt{\frac{[n][N+n]}{[n+1][N+n-1]}} & \sqrt{\frac{[N-1]}{[4][N+2]}} \end{array}\right),$$

$$a_{ts}\begin{bmatrix} R_1 & R_2 \\ \bar{R}_2 & \bar{R}_1 \end{bmatrix} = \frac{1}{K}\left(\begin{array}{c|cc} & s = 0 & \ytableaushort{{\cdot}{}} \\ \hline t = \overset{\longleftarrow n \longrightarrow}{\ydiagram{5,1}} & -\sqrt{\frac{[N-1][N]\dots[N+n-1]}{[n+1][n-1]!}} & \sqrt{\frac{[N][N+1]\dots[N+n]}{[n+1]!}} \\ \overset{\longleftarrow n+1 \longrightarrow}{\ydiagram{6}} & \sqrt{\frac{[N][N+1]\dots[N+n]}{[n+1]!}} & \sqrt{\frac{[N-1][N]\dots[N+n-1]}{[n+1][n-1]!}} \end{array}\right),$$

where $K = \sqrt{dim_q R_1 dim_q R_2}$.

4. Similar exercise could be done for $R_1 = \ydiagram{1}$, $R_2 = \ydiagram{1,1}$, and their conjugate representations are $\bar{R}_1 = \ytableaushort{{\cdot}}$, $\bar{R}_2 = \ydiagram{1,1,1,1}\updownarrow N-2$. These representations give

$$a_{ts}\begin{bmatrix} R_1 & \bar{R}_1 \\ R_2 & \bar{R}_2 \end{bmatrix} = \frac{1}{\sqrt{dim_q R_1\, dim_q R_2}}\left(\begin{array}{c|cc} & s = \ytableaushort{{}{},{\cdot}} & \ytableaushort{{\cdot}{},\none{}} \\ \hline t = 0 & \sqrt{[N]} & \sqrt{\frac{[N-2][N][N+1]}{[2]}} \\ \ytableaushort{{\cdot}{}} & \sqrt{\frac{[N-2][N][N+1]}{[2]}} & -\sqrt{[N]} \end{array}\right),$$

$$a_{ts}\begin{bmatrix} R_1 = \ydiagram{1} & R_2 = \ydiagram{1,1} \\ \bar{R}_1 = \ytableaushort{{\cdot}} & \bar{R}_2 = \ydiagram{1,1,1,1}\updownarrow N-2 \end{bmatrix} = \left(\begin{array}{c|cc} & s = \ytableaushort{{}{},{\cdot}} & \ytableaushort{{\cdot}{},\none{}} \\ \hline t = \ydiagram{2,1} & \sqrt{\frac{[N+1]}{[3][N-1]}} & -\sqrt{\frac{[2][N-2]}{[3][N-1]}} \\ \ydiagram{1,1,1} & -\sqrt{\frac{[2][N-2]}{[3][N-1]}} & -\sqrt{\frac{[N+1]}{[3][N-1]}} \end{array}\right),$$

$$a_{ts}\begin{bmatrix} R_1 & R_2 \\ \bar{R}_2 & \bar{R}_1 \end{bmatrix} = \frac{1}{\sqrt{dim_q R_1\, dim_q R_2}}\left(\begin{array}{c|cc} & s = 0 & \ytableaushort{{\cdot}{}} \\ \hline t = \ydiagram{2,1} & \sqrt{\frac{[N-1][N][N+1]}{[3]}} & -\sqrt{\frac{[N-2][N][N-1]}{[2][3]}} \\ \ydiagram{1,1,1} & -\sqrt{\frac{[N-2][N][N-1]}{[2][3]}} & -\sqrt{\frac{[N-1][N][N+1]}{[3]}} \end{array}\right).$$

5. These results can be generalised for $R_1 = \square$ and $R_2 = \left.\begin{array}{|c|}\hline \\\hline \\\hline \\\hline\end{array}\right\updownarrow n$, whose conjugate representations are $\bar{R}_1 = \boxdot$, $\bar{R}_2 = \left.\begin{array}{|c|}\hline \\\hline \\\hline \\\hline\end{array}\right\updownarrow N-n$ as follows:

$$a_{ts}\begin{bmatrix} R_1 & \bar{R}_1 \\ R_2 & \bar{R}_2 \end{bmatrix} = \frac{1}{K}\left(\begin{array}{c|cc} & s = \left.\begin{array}{|c|}\hline \\\hline \\\hline \\\hline\end{array}\right\updownarrow n-1 & \left.\begin{array}{|c|c|}\hline \cdot & \\\hline \multicolumn{1}{c|}{} & \\\cline{2-2} \multicolumn{1}{c|}{} & \\\cline{2-2} \multicolumn{1}{c|}{} & \\\cline{2-2}\end{array}\right\updownarrow n \\ \hline t=0 & \sqrt{\frac{[N][N-1]...[N-n+2]}{[n-1]!}} & \sqrt{\frac{[N+1][N]...[N-n]}{[N-n+1][n]!}} \\ \begin{array}{|c|c|}\hline \cdot & \\\hline\end{array} & \sqrt{\frac{[N+1][N]...[N-n]}{[N-n+1][n]!}} & -\sqrt{\frac{[N][N-1]...[N-n+2]}{[n-1]!}} \end{array}\right),$$

$$a_{ts}\begin{bmatrix} R_1 & R_2 \\ \bar{R}_1 & \bar{R}_2 \end{bmatrix} = \left(\begin{array}{c|cc} & s = \left.\begin{array}{|c|}\hline \\\hline \\\hline \\\hline\end{array}\right\updownarrow n-1 & \left.\begin{array}{|c|c|}\hline \cdot & \\\hline \multicolumn{1}{c|}{} & \\\cline{2-2} \multicolumn{1}{c|}{} & \\\cline{2-2} \multicolumn{1}{c|}{} & \\\cline{2-2}\end{array}\right\updownarrow n \\ \hline t = \begin{array}{|c|c|}\hline & \\\hline & \multicolumn{1}{c}{}\\\cline{1-1} & \multicolumn{1}{c}{}\\\cline{1-1} & \multicolumn{1}{c}{}\\\cline{1-1}\end{array}\updownarrow n-1 & \sqrt{\frac{[N+1]}{[N+n][N-n+1]}} & -\sqrt{\frac{[n][N-n]}{[n+1][N-n+1]}} \\ \left.\begin{array}{|c|}\hline \\\hline \\\hline \\\hline \\\hline\end{array}\right\updownarrow n+1 & -\sqrt{\frac{[n][N-n]}{[n+1][N-n+1]}} & -\sqrt{\frac{[n][N-n]}{[n+1][N-n+1]}} \end{array}\right),$$

$$a_{ts}\begin{bmatrix} R_1 & R_2 \\ \bar{R}_2 & \bar{R}_1 \end{bmatrix} = \frac{(-1)^n}{K}\left(\begin{array}{c|cc} & s=0 & \begin{array}{|c|c|}\hline \cdot & \\\hline\end{array} \\ \hline t = \begin{array}{|c|c|}\hline & \\\hline & \multicolumn{1}{c}{}\\\cline{1-1} & \multicolumn{1}{c}{}\\\cline{1-1} & \multicolumn{1}{c}{}\\\cline{1-1}\end{array}\updownarrow n-1 & \sqrt{\frac{[N+1][N]\ldots[N-n+1]}{[n+1][n-1]!}} & -\sqrt{\frac{[N][N-1]\ldots[N-n]}{[n]!}} \\ \left.\begin{array}{|c|}\hline \\\hline \\\hline \\\hline \\\hline\end{array}\right\updownarrow n+1 & -\sqrt{\frac{[N][N-1]\ldots[N-n]}{[n]!}} & -\sqrt{\frac{[N+1][N]\ldots[N-n+1]}{[n+1][n-1]!}} \end{array}\right),$$

where $K = \sqrt{dim_q R_1 dim_q R_2}$.

4.2 Coefficients for R_1 symmetric or antisymmetric representations

1. For $R = \begin{array}{|c|c|}\hline & \\\hline\end{array}$, $\bar{R} = \begin{array}{|c|c|}\hline \cdot & \cdot\\\hline\end{array}$

$$a_{ts}\begin{bmatrix} R & \bar{R} \\ R & \bar{R} \end{bmatrix} = \frac{1}{K}\left(\begin{array}{c|ccc} & \overset{\tilde{\rho}_0}{s=0} & \overset{\tilde{\rho}_1}{\begin{array}{|c|c|}\hline \cdot & \\ \hline\end{array}} & \overset{\tilde{\rho}_2}{\begin{array}{|c|c|c|c|}\hline \cdot & \cdot & & \\ \hline\end{array}} \\ \hline t=\tilde{\rho}_0 & \sqrt{dim_q\tilde{\rho}_0} & -\sqrt{dim_q\tilde{\rho}_1} & \sqrt{dim_q\tilde{\rho}_2} \\ \tilde{\rho}_1 & -\sqrt{dim_q\tilde{\rho}_1} & \frac{dim_q\tilde{\rho}_2}{dim_qR-1}-1 & \frac{\sqrt{dim_q\tilde{\rho}_1 dim_q\tilde{\rho}_2}}{dim_qR-1} \\ \tilde{\rho}_2 & \sqrt{dim_q\tilde{\rho}_2} & \frac{\sqrt{dim_q\tilde{\rho}_1 dim_q\tilde{\rho}_2}}{dim_qR-1} & \frac{dim_q\tilde{\rho}_1}{dim_qR-1}-1 \end{array}\right),$$

where

$$K = dim_qR = \frac{[N][N+1]}{[2]},\ dim_q\tilde{\rho}_0 = 1,\ dim_q\tilde{\rho}_1 = [N+1][N-1],$$
$$dim_q\tilde{\rho}_2 = \frac{[N-1][N]^2[N+3]}{[2]^2}\ . \tag{4.3}$$

The second type Racah matrix coefficients are

$$a_{ts}\begin{bmatrix} R & R \\ \bar{R} & \bar{R} \end{bmatrix} = \frac{1}{K}\left(\begin{array}{c|ccc} \text{t} & s=\overset{\tilde{\rho}_0}{0} & \overset{\tilde{\rho}_1}{\begin{array}{|c|c|}\hline \cdot & \\ \hline\end{array}} & \overset{\tilde{\rho}_2}{\begin{array}{|c|c|c|c|}\hline \cdot & \cdot & & \\ \hline\end{array}} \\ \hline \overset{(\rho_1)}{\begin{array}{|c|c|}\hline & \\ \hline & \\ \hline\end{array}} & \sqrt{dim_q\rho_1} & -\frac{[N][N+1]}{[2]}\sqrt{\frac{1}{[3]}} & \frac{[N]}{[2]}\sqrt{\frac{[N+3][N+1]}{[3]}} \\ \overset{(\rho_2)}{\begin{array}{|c|c|c|}\hline & & \\ \hline \multicolumn{1}{|c|}{} \\ \cline{1-1}\end{array}} & -\sqrt{dim_q\rho_2} & x\sqrt{\frac{[N][N+2]}{[4][2]}} & y\sqrt{\frac{[N][N+1][N+2][N+3]}{[4][2]}} \\ \overset{(\rho_3)}{\begin{array}{|c|c|c|c|}\hline & & & \\ \hline\end{array}} & \sqrt{dim_q\rho_3} & u\sqrt{\frac{[N-1][N][N+2][N+3]}{[4][3][2]}} & v\sqrt{\frac{[N-1][N][N+1][N+2]}{[4][3][2]}} \end{array}\right),$$

where the quantum dimensions of the representations are

$$dim_q\rho_1 = \frac{[N-1][N]^2[N+1]}{[2]^2[3]},\ dim_q\rho_2 = \frac{[N-1][N][N+1][N+2]}{[4][2]}\ ,$$
$$dim_q\rho_3 = \frac{[N][N+1][N+2][N+3]}{[4][3][2]}\ . \tag{4.4}$$

The variables x, y, u, v are

$$x = \frac{[N+3][N]}{[N+2]} - [N-1],\ y = \frac{[N]}{[N+2]},\ u = \frac{[2][N+1]}{[N+2]},\ v = \frac{[N+1][2]}{[N+2]} - 1\ .$$

2. For $R=$, $\bar{R}=$ 

$$a_{ts}\left[\begin{array}{cc} R & \bar{R} \\ R & \bar{R} \end{array}\right]=\frac{1}{K}$$

	$s=\overset{\tilde{\rho}_0}{0}$	$\tilde{\rho}_1$ 	$\tilde{\rho}_2$ 
$t=\tilde{\rho}_0$	$\sqrt{dim_q\tilde{\rho}_0}$	$-\sqrt{dim_q\tilde{\rho}_1}$	$\sqrt{dim_q\tilde{\rho}_2}$
$\tilde{\rho}_1$	$-\sqrt{dim_q\tilde{\rho}_1}$	$\frac{dim_q\tilde{\rho}_2}{dim_qR-1}-1$	$\frac{\sqrt{dim_q\tilde{\rho}_1 dim_q\tilde{\rho}_2}}{dim_qR-1}$
$\tilde{\rho}_2$	$\sqrt{dim_q\tilde{\rho}_2}$	$\frac{\sqrt{dim_q\tilde{\rho}_1 dim_q\tilde{\rho}_2}}{dim_qR-1}$	$\frac{dim_q\tilde{\rho}_1}{dim_qR-1}-1$

,

where

$$K=dim_qR=\frac{[N][N-1]}{[2]},\ dim_q\tilde{\rho}_0=1,\ dim_q\tilde{\rho}_1=[N+1][N-1],$$
$$dim_q\tilde{\rho}_2=\frac{[N+1][N]^2[N-3]}{[2]^2}\ . \qquad (4.5)$$

The second type Racah matrix coefficients are

$$a_{ts}\left[\begin{array}{cc} R & R \\ \bar{R} & \bar{R} \end{array}\right]=\frac{1}{K}$$

t	$s=\overset{\tilde{\rho}_0}{0}$	$\tilde{\rho}_1$	$\tilde{\rho}_2$ 
(ρ_1)	$\sqrt{dim_q\rho_1}$	$\frac{[N][N-1]}{[2]}\sqrt{\frac{1}{[3]}}$	$\frac{[N]}{[2]}\sqrt{\frac{[N-3][N-1]}{[3]}}$
 (ρ_2)	$-\sqrt{dim_q\rho_2}$	$x\sqrt{\frac{[N][N-2]}{[4][2]}}$	$y\sqrt{\frac{[N][N-1][N-2][N-3]}{[4][2]}}$
 (ρ_3)	$\sqrt{dim_q\rho_3}$	$-u\sqrt{\frac{[N+1][N][N-2][N-3]}{[4][3][2]}}$	$v\sqrt{\frac{[N+1][N][N-1][N-2]}{[4][3][2]}}$

,

where

$$\begin{aligned} dim_q\rho_1 &= \frac{[N-1][N]^2[N+1]}{[2]^2[3]}, \\ dim_q\rho_2 &= \frac{[N+1][N][N-1][N-2]}{[4][2]}, \\ dim_q\rho_3 &= \frac{[N][N-1][N-2][N-3]}{[4][3][2]}\ , \end{aligned}$$

and the variables are

$$x=[N+1]-\frac{[N-3][N]}{[N-2]},\qquad y=\frac{[N]}{[N-2]},$$
$$u=\frac{[2][N-1]}{[N-2]},\qquad v=\frac{[N-1][2]}{[N-2]}-1\ . \tag{4.6}$$

3. For $R_1=\Box\Box\neq R_2=\Box\Box\Box$, which is will be useful for the computation of links, the second type Racah coefficients are

$$a_{ts}\begin{bmatrix} R_1 & R_2\\ \bar{R}_1 & \bar{R}_2\end{bmatrix}=\left(\begin{array}{c|ccc} \text{t} & s=\overset{\tilde{\rho}_0}{\Box} & \overset{\tilde{\rho}_1}{\boxdot\Box\Box} & \overset{\tilde{\rho}_2}{\boxdot\boxdot\Box\Box\Box} \\ \hline \rho_0 & z_1\sqrt{\frac{[N-1][N][N+1][N+2]}{[4][3][2]}} & -\frac{[2]}{[N+1]}\sqrt{\frac{[N][N+1]}{[4][3]}} & \frac{[2]}{[N+2]}\sqrt{\frac{[N+2][N+4]}{[4][2]}} \\ \rho_1 & -z_1[2]\sqrt{\frac{[N-1][N+1][N+2][N+3]}{[2][3][5]}} & z_2\sqrt{\frac{[N+1][N+3]}{[3][5]}} & z_3\sqrt{\frac{[N][N+2][N+3][N+4]}{[2][5]}} \\ \rho_2 & z_1[3]\sqrt{\frac{[N+1][N+2][N+3][N+4]}{[5][4][3][2]}} & z_3\sqrt{\frac{[N-1][N+1][N+3][N+4]}{[5][4][3]}} & z_2\sqrt{\frac{[N-1][N][N+2][N+3]}{[5][4][2]}} \end{array}\right),$$

where $\rho_0=\begin{smallmatrix}\Box\Box\Box\\ \Box\Box\end{smallmatrix}$, $\rho_1=\begin{smallmatrix}\Box\Box\Box\Box\\ \Box\end{smallmatrix}$, $\rho_2=\Box\Box\Box\Box\Box$ and the variables z_i's are

$$z_1=\frac{[2]}{[N+1][N+2]},\qquad z_2=z_1\left(\frac{[N+3][N+4]-\frac{[N][N-1]}{[3]}}{\frac{[N]}{[3]}+[N+4]}\right),$$
$$z_3=\frac{[2]}{[N+2][N+3]},\qquad z_4=\frac{[2]}{[N+1][N+3]}\ ,$$

and gets more cumbersome for the third type Racah-matrix:

$$a_{ts}\begin{bmatrix} R_1 & R_2\\ \bar{R}_2 & \bar{R}_1\end{bmatrix}=\frac{1}{dim_qR_1dim_qR_2}\left(\begin{array}{c|ccc} & s=\overset{\tilde{\rho}_0}{0} & \overset{\tilde{\rho}_1}{\boxdot\Box} & \overset{\tilde{\rho}_2}{\boxdot\boxdot\Box\Box} \\ \hline t=\rho_0 & \sqrt{dim_q\rho_0} & -[N]\sqrt{\frac{[N+3]}{[4]}} & [N]\sqrt{\frac{[N-1][N+3][N+4]}{[2][3][4]}} \\ \rho_1 & -\sqrt{dim_q\rho_1} & x\sqrt{\frac{[N]}{[5]}} & y\sqrt{\frac{[N-1][N][N+4]}{[2][3][5]}} \\ \rho_2 & \sqrt{dim_q\rho_2} & u\sqrt{\frac{[N-1][N][N+4]}{[4][5]}} & v\sqrt{\frac{[N]}{[2][3][4][5]}} \end{array}\right).$$

The quantum dimensions of the representations in terms of the q-numbers are

$$\sqrt{dim_qR_1dim_qR_2}=\frac{[N][N+1]}{[2]}\sqrt{\frac{[N+2]}{[3]}},\ dim_q\rho_0=\frac{[N-1][N]^2[N+1][N+2]}{[4][3][2]},$$
$$dim_q\rho_1=\frac{[N-1][N][N+1][N+2][N+3]}{[2][3][5]},\ dim_q\rho_2=\frac{[N][N+1][N+2][N+3][N+4]}{[2][3][4][5]}$$

and the variables are

$$x = \frac{[N-1]}{[2][3]}\left(y[N+4]-[N+1][N+2]\right), \qquad y = \frac{b_1+\sqrt{b_1^2-a_1c_1}}{a_1}$$

$$a_1 = \frac{[N-1][N+4]}{[2][3]} + \left(\frac{[N-1][N+4]}{[2][3]}\right)^2, \qquad b_1 = \left(\frac{[N-1]}{[2][3]}\right)^2[N+1][N+2][N+4]$$

$$u = \frac{[N-1]}{[2][3]}\left(y[N+4]-[N+1][N+2]\right), \qquad v = 1+\sqrt{1-\frac{c_2}{a_2}},$$

$$\begin{aligned}
a_2 &= [N][2][3]\left([N-1][N+4]+[2][3]\right),\\
c_1 &= \frac{[N-1][N+1][N+2][N+3]}{[2][3]} + \left(\frac{[N-1][N+1][N+2]}{[2][3]}\right)^2\\
&\quad -\frac{[5][N+2]}{[3][N]}\left(\frac{[N][N+1]}{[2]}\right)^2,\\
c_2 &= [N][N+1][N+2][N+3][N+4]+[N][N+1]^2[N+2]^2\\
&\quad -\frac{[N]^2[N+1]^2}{[2]}[N+2][4][5]\ .
\end{aligned}$$

Using the identities, it should be possible to generalise these Racah matrices for $R_2 = \overset{\xleftrightarrow{n}}{\square\square\square\square\square\square}$, which are again 3×3 matrices.

4. Equivalently, we could write the Racah matrix coefficients when $R_1 = \begin{smallmatrix}\square\\\square\end{smallmatrix} \neq R_2$ where R_2 is totally antisymmetric n-th rank tensor (represented by n-vertical box). For $R_1 = \begin{smallmatrix}\square\\\square\end{smallmatrix}$ and $R_2 = \begin{smallmatrix}\square\\\square\\\square\end{smallmatrix}$, the second type Racah coefficient matrix is

$$a_{ts}\begin{bmatrix} R_1 & R_2\\ \bar{R}_1 & \bar{R}_2\end{bmatrix} = \left(\begin{array}{c|ccc}
\text{t} & s=\overset{\tilde\rho_0}{\square} & \overset{\tilde\rho_2}{\begin{smallmatrix}\square&\square\\&\square\end{smallmatrix}} & \overset{\tilde\rho_2}{N-4\updownarrow\begin{smallmatrix}\square&\square\\\square&\square\\\square&\square\\\square\end{smallmatrix}}\\
\hline
\rho_0 & z_1\sqrt{\frac{[N+1][N][N-1][N-2]}{[4][3][2]}} & \frac{[2]}{[N-1]}\sqrt{\frac{[N][N-1]}{[4][3]}} & \frac{[2]}{[N-2]}\sqrt{\frac{[N-2][N-4]}{[4][2]}}\\
\rho_1 & -z_1[2]\sqrt{\frac{[N+1][N-1][N-2][N-3]}{[2][3][5]}} & z_2\sqrt{\frac{[N-1][N-3]}{[3][5]}} & z_3\sqrt{\frac{[N][N-2][N-3][N-4]}{[2][5]}}\\
\rho_2 & z_1[3]\sqrt{\frac{[N-1][N-2][N-3][N-4]}{[5][4][3][2]}} & -z_3\sqrt{\frac{[N+1][N-1][N-3][N-4]}{[5][4][3]}} & z_2\sqrt{\frac{[N+1][N][N-2][N-3]}{[5][4][2]}}
\end{array}\right),$$

where $\rho_0 = \begin{smallmatrix}\square&\square\\\square&\square\\\square\end{smallmatrix}$, $\rho_1 = \begin{smallmatrix}\square&\square\\\square\\\square\\\square\end{smallmatrix}$, $\rho_2 = \begin{smallmatrix}\square\\\square\\\square\\\square\\\square\end{smallmatrix}$ and the variables are

$$z_1 = \frac{[2]}{[N-1][N-2]}, \qquad z_2 = a\left(\frac{-[N-3][N-4]+\frac{[N][N+1]}{[3]}}{\frac{[N]}{[3]}+[N-4]}\right),$$

$$z_3 = \frac{[2]}{[N-2][N-3]}, \qquad z_4 = \frac{[2]}{[N-1][N-3]}\ .$$

Finally, with these data available, we can evaluate the polynomial form for the non-torus knots and links in Figs. 1&2.

5 Computation of non-torus link invariants

Now with the Racah coefficients for some representations determined in the previous section, we will now explicitly work out the polynomial invariant for a non-torus framed knot and a framed two component link. From these invariants, the reformulated invariants $f_R[K], f_{R_1,R_2}[L]$ (2.22) are obtained and shown to obey Ooguri-Vafa conjecture (2.24).

5.1 Non-torus knot 5_2

Using eqn.(2.19), the $U(N)$ invariants for the zero-framed knot 5_2 $((p+p_1)=0)$ carrying representation $\square$, $\square\!\square$ and $\begin{smallmatrix}\square\\[-2pt]\square\end{smallmatrix}$ (2.20) are

$$V_{\square}^{\{U(N)\}}[5_2;0] \equiv W_{\square}(q,\lambda) = \frac{1}{q^{\frac{1}{2}}-q^{-\frac{1}{2}}}[q^{-1}\lambda^{-7/2}(q-\lambda+\lambda^3-q^2\lambda-q\lambda^3+q^2\lambda^3)] \tag{5.1}$$

$$\begin{aligned} V_{\square\!\square}^{\{U(N)\}}[5_2;0] &\equiv W_{\square\!\square}(q,\lambda) \\ &= \frac{1}{(q-1)^2(q+1)}[q^{-5}\lambda^{-7}(q-\lambda+2\lambda^2-\lambda^3+(q\lambda)^5+2(q\lambda)^6 \\ &\quad -q^8\lambda^4-2q\lambda-q^4\lambda+q\lambda^2+q^3\lambda^2+q^4\lambda^2-q^5\lambda^2-q^3\lambda^3+q^4\lambda^3 \\ &\quad +q^5\lambda^3+q^7\lambda^3+q^8\lambda^3-q^5\lambda^4-q^6\lambda^4-q^9\lambda^4-q^3\lambda^5-q^6\lambda^5 \\ &\quad +q^3\lambda^6-q^4\lambda^6-q^7\lambda^6+q^8\lambda^5-q^8\lambda^6+q^9\lambda^6)]\ , \end{aligned} \tag{5.2}$$

$$\begin{aligned} V_{\begin{smallmatrix}\square\\[-2pt]\square\end{smallmatrix}}^{\{U(N)\}}[5_2;0] &\equiv W_{\begin{smallmatrix}\square\\[-2pt]\square\end{smallmatrix}}(q,\lambda) \\ &= \frac{1}{(q-1)^2(q+1)}[q^{-1}\lambda^{-7}(q^8-\lambda^4+\lambda^6+(q\lambda)^6-q^6\lambda^3+q^6\lambda^2 \\ &\quad -q^9\lambda^3+q^8\lambda^2-q^5\lambda-2q^8\lambda-q^9\lambda-q^4\lambda^2+q^5\lambda^2+2q^9\lambda^2 \\ &\quad q\lambda^3+q^2\lambda^3+q^4\lambda^3+q^5\lambda^3-q\lambda^4-q^3\lambda^4-q^4\lambda^4+q\lambda^5-q^3\lambda^5 \\ &\quad q^4\lambda^5-q^6\lambda^5-q\lambda^6-q^2\lambda^6+2q^3\lambda^6-q^5\lambda^6)]\ . \end{aligned} \tag{5.3}$$

Using the relation between the reformulated invariants in terms of the Chern-Simons invariants (2.22), $f_{\square\square}$ and $f_{\begin{smallmatrix}\square\\\square\end{smallmatrix}}$ will be

$$f_{\square\square}(q,\lambda) = W_{\square\square}(q,\lambda) - \frac{1}{2}\left(W_{\square}(q,\lambda)^2 + W_{\square}(q^2,\lambda^2)\right) \ , \tag{5.4}$$

$$f_{\begin{smallmatrix}\square\\\square\end{smallmatrix}}(q,\lambda) = W_{\begin{smallmatrix}\square\\\square\end{smallmatrix}}(q,\lambda) - \frac{1}{2}\left(W_{\square}(q,\lambda)^2 - W_{\square}(q^2,\lambda^2)\right) \ . \tag{5.5}$$

Substituting the $U(N)$ invariants for 5_2 knot, we find

$$\begin{aligned} f_{\square\square}(q,\lambda) &= \frac{1}{q^{\frac{1}{2}} - q^{-\frac{1}{2}}}[q^{-\frac{11}{2}}\lambda^{-7}(1-q+q^2)(-1+\lambda)^2(q+q^2+q^3-\lambda \\ &\quad -q\lambda - q^2\lambda - q^4\lambda + q^3\lambda^2 - q^4\lambda^2 + q^3\lambda^3 - q^4\lambda^3 + q^3\lambda^4)] \ , \end{aligned} \tag{5.6}$$

$$\begin{aligned} f_{\begin{smallmatrix}\square\\\square\end{smallmatrix}}(q,\lambda) &= \frac{1}{q^{\frac{1}{2}} - q^{-\frac{1}{2}}}[q^{-\frac{1}{2}}\lambda^{-7}(1-q+q^2)(-1+\lambda)^2(q+q^2+q^3-\lambda \\ &\quad -\lambda^2 - \lambda^3 - q^2\lambda - q^3\lambda - q^4\lambda + q\lambda^2 + q\lambda^3 + q\lambda^4)] \ , \end{aligned} \tag{5.7}$$

obeying the Ooguri-Vafa conjecture(2.24). Conversely, this confirms the correctness of the second type Racah coefficients tabulated in the previous section.

5.2 Non-torus 6_2 link

We can again compute the invariant for the link 6_2 (2.21) carrying same or different representations and framing $p_1 = p_2 = 0$ on the component knots:

$$\begin{aligned} V^{\{U(N)\}}_{(\square,\square)}[6_2;0] &\equiv W_{(\square,\square)}(q,\lambda) \\ &= \frac{1}{(q-1)^2}(q - q^{-1} - 2q + q^2 - q^3 - q\lambda^{-1} + q^2\lambda^{-1} + \lambda^{-1} - 2q^2 \\ &\quad +q\lambda + q^{-1}\lambda^2 + 2q\lambda^2 - 2q^2\lambda^2 + q^2\lambda^2) \ , \end{aligned} \tag{5.8}$$

$$\begin{aligned} V^{\{U(N)\}}_{(\square\square,\square)}[6_2;0] &\equiv W_{(\square\square,\square)}(q,\lambda) \\ &= \frac{1}{(q-1)^3(q+1)}[q^{-\frac{1}{2}}\lambda^{-\frac{3}{2}}(-q+q^3-q^4+\lambda-\lambda^2+q^8\lambda^4-q^5\lambda \\ &\quad +q^6\lambda + q\lambda - q^2\lambda + q^3\lambda + q^4\lambda - q^3\lambda^2 - q^5\lambda^2 + q^4\lambda^3 + q^7\lambda^3 \\ &\quad -q^8\lambda^3 + q^2\lambda^4 - q^3\lambda^4 - q^4\lambda^4 + 2q^5\lambda^4 - q^6\lambda^4 - q^7\lambda^4)] \ , \end{aligned} \tag{5.9}$$

$$\begin{aligned} V^{\{U(N)\}}_{(\begin{smallmatrix}\Box\\\Box\end{smallmatrix},\Box)}[6_2;0] &\equiv W_{(\begin{smallmatrix}\Box\\\Box\end{smallmatrix},\Box)}(q,\lambda) \\ &= \frac{1}{(q-1)^3(q+1)}[q^{-\frac{7}{2}}\lambda^{-\frac{3}{2}}(-q^4+q^5-q^7-\lambda^3+\lambda^4-q^8\lambda^2 \\ &\quad q^5\lambda-q^6\lambda+q^7\lambda+q^8\lambda+q^2\lambda-q^3\lambda+q^4\lambda-q^3\lambda^2-q^5\lambda^2 \\ &\quad +q\lambda^3+q^4\lambda^3-q\lambda^4-q^2\lambda^4+2q^3\lambda^4-q^4\lambda^4-q^5\lambda^4 \\ &\quad +q^6\lambda^4)]\ . \end{aligned} \tag{5.10}$$

Following eqn.(2.22), the reformulated invariant for two-component link simplifies as

$$\begin{aligned} f_{(\Box\!\Box,\Box)}(\mathcal{L}) &= W_{(\Box\!\Box,\Box)}(\mathcal{L})-W_{(\Box,\Box)}(\mathcal{L})W_{\Box}(\mathcal{K}_1)-W_{\Box\!\Box}(\mathcal{K}_1)W_{\Box}(\mathcal{K}_2)+W_{\Box}(\mathcal{K}_1)^2W_{\Box}(\mathcal{K}_2) \\ f_{(\begin{smallmatrix}\Box\\\Box\end{smallmatrix},\Box)}(\mathcal{L}) &= W_{(\begin{smallmatrix}\Box\\\Box\end{smallmatrix},\Box)}(\mathcal{L})-W_{(\Box,\Box)}(\mathcal{L})W_{\Box}(\mathcal{K}_1)-W_{\begin{smallmatrix}\Box\\\Box\end{smallmatrix}}(\mathcal{K}_1)W_{\Box}(\mathcal{K}_2)+W_{\Box}(\mathcal{K}_1)^2W_{\Box}(\mathcal{K}_2). \end{aligned}$$

Again, we get the expected Ooguri-Vafa conjectured form[7] for 6_2 link (2.24):

$$\begin{aligned} f_{(\Box\!\Box,\Box)}(q,\lambda) &= (q\lambda)^{-\frac{1}{2}}(q^2-\lambda^2+\lambda^3-2q^2\lambda^2-q^3-q^4\lambda^2+q\lambda^3\lambda^2 \\ &\quad +q^2\lambda^3+q^3\lambda^3+q^4\lambda^3)\ , \qquad (5.11) \\ f_{(\begin{smallmatrix}\Box\\\Box\end{smallmatrix},\Box)}(q,\lambda) &= -(q^{-\frac{7}{2}}\lambda^{-\frac{1}{2}})(q^2-\lambda^2+\lambda^3-q\lambda^2-2q^2\lambda^2-q^3\lambda^2 \\ &\quad -q^4\lambda^2+q\lambda^3+q^2\lambda^3+q^3\lambda^3+q^4\lambda^3)\ . \qquad (5.12) \end{aligned}$$

6 Discussion and Conclusion

In this paper, we have attempted a challenging problem of obtaining matrix elements of the duality matrix which has properties and identities similar to the quantum Racah coefficients. Particularly, we derived these identities and properties by studying the equivalence of states in the space of correlator conformal blocks in the $SU(N)_k$ Wess-Zumino conformal field theory.

We have tabulated the Racah coefficients for some class of representations which will be useful to compute non-torus knots and non-torus two component links. We verified the polynomial form of the 5_2 knot and 6_2 link and obtained their reformulated invariants. These invariants obey the form conjectured by Ooguri-Vafa[6, 7] confirming the correctness of our Racah coefficients in section 4.

We believe that there must be a systematic way of writing a closed form expression similar to the expression obtained for $SU(2)$ quantum Racah coefficients

[5, 16]. There are papers in the literature addressing classical Racah and quantum Racah coefficients. Unfortunately, we do not see such explicit coefficients in section 4 to compare. We hope to study those papers which may help us to obtain a closed form expression for $SU(N)$ quantum Racah coefficients.

There are interesting recent developments relating torus knots to spectral curve in the B-model topological strings[10], Poincare polynomial computation from refined Chern-Simons theory, Khovanov homology, fivebranes[9, 11], and the polynomial invariants from counting of solutions in four-dimensional theories[12]. We hope to extend these recent works to non-torus links and report in future.

Acknowledgments: P.R would like to thank the hospitality of Center for Quantum spacetime, Sogang University where this work was completed. This work was supported by the National Research Foundation of Korea(NRF) grant funded by the Korea government(MEST) through the Center for Quantum Spacetime(CQUeST) of Sogang University with grant number 2005-0049409.

References

[1] E. Witten, " Quantum field theory and the Jones polynomial," Commun. Math. Phys. **121** (1989), 351.

[2] P. Ramadevi, T.R. Govindarajan, R. K. Kaul, "Three-Dimensional Chern-Simons Theory as a Theory of Knots and Links (III). Compact Semisimple Group," Nucl. Phys. **B 402** (1993) 548, `hep-th/9212110`.

[3] P. Ramadevi, " Chern-Simons theory as a theory of knots and links,", Ph.D. Thesis.

[4] J. M. F. Labastida, P. M. Llatas, A. V. Ramallo, "Knot Operators in Chern-Simons theory," Nucl. Phys. **B348** (1991) 651.

[5] A.N. Kirillov, N.Yu.Reshetikhin,"Representation algebra $U_q(SL(2))$, q-orthogonal polynomials and invariants of links. LOMI preprint E-9-88; see also in: New Developments in the theory of knots. ed. T. Kohno,Singapore: World Scientific, 1989.

[6] H. Ooguri, C. Vafa, "Knot Invariants and Topological Strings," Nucl. Phys. **B577**, 419, (2000), `hep-th/9912123`.

[7] J.M.F. Labastida, M. Marino, C. Vafa, " Knots, links and branes at large N,"JHEP 0011:007, (2000), `hep-th/0010102`.

[8] P. Ramadevi, T. Sarkar, "On Link Invariants and Topological String Amplitudes," Nucl. Phys. **B 600** (2001) 487.

[9] E. Witten, "Fivebranes and knots," arXiv:1101.3216 [hep-th].

[10] A. Brini, B. Eynard and M. Marino,"Torus Knots and mirror symmetry," arXiv:1105.2012[hep-th]

[11] M. Aganagic, S. Shakirov,"Knot homology from refined Chern-Simons theory," arXiv:1105.5117[hep-th]

[12] D. Gaitto, E. Witten, "Knot Invariants from Four-Dimensional Gauge Theory," arXiv:1106.4789 [hep-th]

[13] M. Marino, C. Vafa, " Framed Knots at Large N," `hep-th/0108064`.

[14] P. Borhade, P. Ramadevi, T. Sarkar,"$U(N)$ Framed Links, Three-Manifold Invariants, and Topological Strings," Nucl.Phys. B678 (2004) 656.

[15] J.S. Birman,"Braids, Links and Mapping Class groups," Annals of Mathematics Studies, Princeton, NJ: Princeton Univ. Press, 1975.

[16] R.K. Kaul,"Chern-Simons Theory,Colored-Oriented Braids and Link Invariants," Commun. Math. Phys. **162** (1994) 289.

[17] J.M.F. Labastida, M.Marino,"A New Point of View in the Theory of Knot and Link Invariants,"J.Knot Theory Ramifications **11** (2002) 173.

[18] Chandrima Paul, Pravina Borhade, P. Ramadevi,"Composite Representation Invariants and Unoriented Topological String Amplitudes," Nucl.Phys.**B841** (2010) 448.